


# Estimating Consequences of 3-Body Forces

Harald W. Griebhammer\*

Center for Nuclear Studies, The George Washington University, Washington DC 20052, USA

**Abstract.** Classifying the strengthes of three-body forces 3BFs with the condition that observables must be cut-off independent, i.e. renormalised at each order, leads to surprising results with relevance for example for thermal neutron capture on the deuteron. Details and a better bibliography in Ref. [1].

Adding 3-Body Forces (3BFs) *a posteriori* when theory and data disagree is untenable when predictions are required. Effective Field Theories (EFTs), see e.g. [2] for reviews, provide a model-independent way to estimate their typical strength. For systems of three identical particles in which short-range forces produce shallow two-particle bound states, and in particular for the “pion-less” EFT of Nuclear Physics EFT( $\not{\pi}$ ), consistency arguments from renormalisation lead to a power-counting, namely a recipe to systematically estimate the typical size of 3BFs in all partial waves and orders, including external currents.



**Figure 1.** Left: integral equation of nucleon-deuteron scattering. Right: generic loop correction (rectangle) at  $N^n$ LO. Thick line ( $D$ ):  $NN$  propagator; thin line ( $\mathcal{K}_l$ ): propagator of the exchanged nucleon; ellipse: LO half off-shell amplitude.

We start from the Faddeev equation, Fig. 1, in the  $l$ th partial wave of the spin doublet ( $\lambda = 1$ ) and quartet ( $\lambda = -\frac{1}{2}$ ) channels of nucleon-deuteron scattering at leading order LO. The  $NN$  amplitude is given by the leading term of the Effective-Range Expansion. The 3-nucleon amplitude converges for large half off-shell momenta  $p$  as  $p^{-s_0-1}$ , with  $s_0$  the solution to the algebraic equation

$$1 = (-1)^l \frac{2^{1-l} \lambda}{\sqrt{3\pi}} \frac{\Gamma\left[\frac{l+s+1}{2}\right] \Gamma\left[\frac{l-s+1}{2}\right]}{\Gamma\left[\frac{2l+3}{2}\right]} {}_2F_1\left[\frac{l+s+1}{2}, \frac{l-s+1}{2}; \frac{2l+3}{2}; \frac{1}{4}\right]. \quad (1)$$

The asymptotics depends thus crucially on  $l$  and the spin-isospin factor  $\lambda$  only.

\*E-mail address: hgrie@gwu.edu; 29 Nov 2007; arXiv:0711.4522. To appear in Few-Body Systems.

As a description in terms of point-like interactions becomes inadequate beyond a maximum momentum, low-energy observables must be insensitive to details of the amplitude at large  $p$ , i.e. to form and value of the regulator chosen. This is the fundamental tenet of EFT: Include a 3BF *if and only if* needed to cancel cut-off dependence in low-energy observables, i.e. as counter-term for divergences which can not be absorbed by renormalising 2-body interactions. 3BFs are thus not added out of phenomenological needs but to guarantee that observables are insensitive to off-shell effects. Counting loop momenta in Fig. 1, the *superficial degree of divergence* of a higher-order correction is non-negative for

$$\text{Re}[n - s_l(\lambda) - s_{l'}(\lambda')] \geq 0 . \quad (2)$$

The order  $n$  at which a 3BF is needed is thus determined just by the order at which a correction to the 3-body amplitude with only 2-body interactions starts to depend on unphysical short-distance behaviour.

Let us re-visit the problem in position space. The Schrödinger equation for the wave-function in the hyper-radial deuteron-nucleon distance  $r$  at short distances,

$$\left[ -\frac{1}{r} \frac{\partial}{\partial r} r \frac{\partial}{\partial r} + \frac{s_l^2(\lambda)}{r^2} - ME \right] F(r) = 0 , \quad (3)$$

looks like the one for a free particle with centrifugal barrier. One would thus expect  $s_l \stackrel{?}{=} l+1$  (hyper-spherical co-ordinates!). It is however well-known that the centrifugal term is attractive for three bosons ( $\lambda = 1$ ), contrary to expectations. The wave-function collapses to the origin and seems infinitely sensitive to very-short-distance physics. In order to stabilise the system – or, equivalently, remove dependence on details of the cut-off –, a 3BF must be added at LO. This leads to the famous “limit-cycle” with its Efimov and Thomas effects and universal correlations, see e.g. Ref. [2] for reviews.

On the other hand, 3BFs are demoted if  $s_l > l + 1$  because the centrifugal barrier provides *more* repulsion than expected. The wave-function is pushed further out and thus *less* sensitive to short-distance details. About half of the 3BFs for  $l \leq 2$  are *weaker*, half *stronger* than one would expect simplistically, see Table 1. The higher partial-waves follow expectation, as the Faddeev equation is then saturated by the Born approximation.

partial-wave: <i>in-out</i>	naïve dim. analysis	simplistic		typ. size
bosons      fermions	$\text{Re}[s_l(\lambda) + s_{l'}(\lambda')]$	$l + l' + 2$		if $Q^n \sim \frac{1}{3^n}$
S-S	LO	$N^2\text{LO}$	promoted	100% (10%)
	$N^{3.1}\text{LO}$	$N^4\text{LO}$	promoted	3% (1%)
P-P	$N^{5.7}\text{LO}$	$N^4\text{LO}$	demoted	0.2% (1%)
	$N^{3.5}\text{LO}$		promoted	2% (1%)
	$N^{4.3+2}\text{LO}$	$N^{2+2}\text{LO}$	demoted	0.1% (1%)
	$N^{5.0}\text{LO}$	$N^4\text{LO}$	demoted	0.4% (1%)
higher	$\sim$ as simplistic	$N^{l+l'+2}\text{LO}$		

**Table 1.** Order of some leading 3BFs in nucleon-deuteron scattering, indicating if actual values from eqs. (1/2) are stronger (“promoted”) or weaker (“demoted”) than the simplistic estimate. Last column: typical size of 3BF in EFT( $\#$ ); in parentheses size from the simplistic estimate.

Demotion might seem an academic dis-advantage – to include some unnecessary higher-order corrections does not improve the accuracy of the result. But demotion is pivotal when predicting the experimental precision necessary to disentangle 3BFs in observables, and here the error-estimate of EFTs is crucial.

An example is the cross-section of triton radiative capture  $nd \rightarrow t\gamma$  at thermal energies, see Ref. [3] for details and references. Nuclear models give a spread of  $[0.49 \dots 0.66]$  mb, depending on the 2-nucleon potential and inclusion of the  $\Delta(1232)$  [4]. Restoring gauge-invariance reduces the spread to  $[0.52 \dots 0.56]$  mb, but the discrepancy to experiment increases when gauge-invariant three-nucleon currents are added [4]. On the other hand, this low-energy process should be insensitive to details of Physics at 300 MeV. Indeed, the power-counting of 3BFs applies equally with external currents. No new 3BFs are needed up to N<sup>2</sup>LO to render cut-off independence. The result converges order by order,

$$\sigma_{\text{tot}} = [0.485(\text{LO}) + 0.011(\text{NLO}) + 0.007(\text{N}^2\text{LO})] \text{ mb} = [0.503 \pm 0.003] \text{ mb} \quad , \quad (4)$$

is cut-off independent and compares well with experiment,  $[0.509 \pm 0.015]$  mb. In contradistinction to earlier potential models, it is manifestly gauge-invariant.

The potential models reproduce the input of EFT( $\not{\neq}$ ): the nucleon magnetic moments, deuteron and triton binding energies,  $NN$  and  $nd$  scattering lengths, and the thermal cross-section of  $np \rightarrow d\gamma$ . That their results vary dramatically is at odds with universality, a key aspect of EFTs: Answers from models with the same input should agree with EFT( $\not{\neq}$ ) within the projected accuracy. This conflict poses a puzzle whose resolution is not (yet) clear.

With these findings, EFT( $\not{\neq}$ ) is a self-consistent field theory which contains the minimal number of interactions at each order to be renormalisable. Each 3-body counter-term gives rise to one subtraction-constant, fixed by a 3-body datum. This method is applicable to any EFT with an infinite number of diagrams at LO, e.g. because of shallow bound-states. It leads at each order and to the prescribed level of accuracy to a cut-off independent theory with the smallest number of experimental input-parameters. The power-counting is thus not constructed by educated guesswork but by investigations of the renormalisation-group properties of couplings and observables using rigorous methodology.

*Acknowledgement.* It is a pleasure to thank the organisers for creating a highly stimulating conference and a warm welcome. Supported in part by the US Department of Energy (DE-FG02-95ER-40907) and a CAREER-grant PHY-0645498 of the US National Science Foundation.

## References

1. H. W. Griesshammer, Nucl. Phys. A **760** (2005) 110.
2. P. F. Bedaque and U. van Kolck, Ann. Rev. Nucl. Part. Sci. **52** (2002) 339; E. Braaten and H. W. Hammer, Phys. Rept. **428** (2006) 259.
3. H. Sadeghi, S. Bayegan, H. W. Griesshammer: Phys. Lett. **B643** (2006) 263.
4. M. Viviani et al., Phys. Rev. **C54** (1996) 534; L. E. Marcucci et al., Phys. Rev. **C72** (2005) 014001.